

Infinitely p -divisible points on abelian varieties defined over function fields of characteristic $p > 0$.

Damian RÖSSLER*

January 14, 2013

Abstract

In this article we consider some questions raised by F. Benoit, E. Bouscaren and A. Pillay. We prove that infinitely p -divisible points on abelian varieties defined over function fields of transcendence degree one over a finite field are necessarily torsion points. We also prove that when the endomorphism ring of the abelian variety is \mathbb{Z} then there are no infinitely p -divisible points of order a power of p .

1 Introduction

Fix once and for all a prime number p .

Let K_0 be the function field of a smooth curve over a finite field \mathbb{F} of characteristic p . Let B be an abelian variety over K_0 .

For any abelian group G , define $G^\# := \cap_{l \geq 0} p^l G$. We call the elements of $G^\#$ the infinitely p -divisible points of G . Furthermore, write $\text{Tor}(G)$ for the subset of G consisting of elements of finite order. If $n \in \mathbb{N}^*$, we write $\text{Tor}^n(G)$ for the subset of $\text{Tor}(G)$ consisting of elements of order prime to n . Similarly, we write $\text{Tor}_n(G)$ for the subset of $\text{Tor}(G)$ consisting of elements whose order divides a power of n .

*Institut de Mathématiques, Equipe Emile Picard, Université Paul Sabatier, 118 Route de Narbonne, 31062 Toulouse cedex 9, FRANCE, E-mail: rossler@math.univ-toulouse.fr

The following conjecture is made at the very end of the article [3].

Conjecture 1.1. *The inclusion $B(K_0^s)^\# \subseteq \text{Tor}(B(K_0^s))$ holds.*

Here K_0^s is the separable closure of K_0 .

In the same context, F. Benoist then asked the following question (see [2]).

Question 1.2. *Suppose that there are no non vanishing \bar{K}_0 -homomorphisms $B_{\bar{K}_0} \rightarrow C$, where C is an abelian variety, which has a model over $\bar{\mathbb{F}}$.*

Do we have $B(K_0^s)^\# \cap \text{Tor}_p(B(K_0^s)) = 0$?

We recall (in the notation of Question 1.2) that a model of C over $\bar{\mathbb{F}}$ is an abelian variety C_0 over $\bar{\mathbb{F}}$ such that $C \simeq C_0 \times_{\bar{\mathbb{F}}} \bar{K}_0$.

In response to these questions, we shall prove the following two results in this text :

Theorem 1.3. *Conjecture 1.1 holds.*

A. Pillay explained that Conjecture 1.1 can be viewed as a positive characteristic analog of Manin's "theorem of the kernel" (see [14] for the latter). He also outlined a proof of the Mordell-Lang conjecture over function fields of characteristic $p > 0$ (in the case of abelian varieties), which is based on Conjecture 1.1 and a quantifier elimination result.

Conjecture 1.1 is also linked to the Mordell-Lang conjecture in the following way. Lemma 2.4 below (which plays a key role in the proof of Conjecture 1.1) implies that the infinitely p -divisible points defined over a certain separable (but transcendental) field extension of K_0 are annulled by a fixed Weil polynomial (in particular it has no roots of unity among its roots) applied to a lifting of the Frobenius automorphism. This fact, combined with the existence of arc schemes (see for instance [23, before Lemma 2.3]) as well as Proposition 6.1 in [19] can be used to give a quick proof of Theorem 4.1 in [23]. This last theorem is the main tool in the proof of the Mordell-Lang conjecture given in [23]; more precisely, the Mordell-Lang conjecture follows quickly from it (see par. 3.2 in [23] for the argument), once the existence of jet schemes (see for instance par. 2.1 in [23] for the latter) is established.

Details about the argument outlined in the last paragraph will appear in [4], where it will also be generalized to the semiabelian situation; it would be very

interesting to understand the link (if there is one) between this argument and A. Pillay's approach to the Mordell-Lang conjecture mentioned above.

Theorem 1.4. *Suppose that $\text{End}_{\bar{K}_0}(B) = \mathbb{Z}$. Then $B(K_0^s)^\# \cap \text{Tor}_p(B(K_0^s)) = 0$.*

Is the answer to F. Benoit's question affirmative if $\text{End}_{\bar{K}_0}(B) = \mathbb{Z}$.

We actually prove a stronger result, but the stronger form is (as very often) more difficult to formulate. See the last remark of the text for this stronger form.

In particular, the answer to Question 1.2 is affirmative if B is an elliptic curve over K_0 . This was also proved in [3]. In [26, after Th. 3], F. Voloch shows that if B/K_0 is ordinary and has maximal Kodaira-Spencer rank, then $B[p](K_0^s) = 0$ and thus the answer to Question 1.2 is also affirmative in that situation.

The hypothesis $B(K_0^s)^\# \cap \text{Tor}_p(B(K_0^s)) = 0$ is a crucial hypothesis in B. Poonen's and J. F. Voloch's work on the Brauer-Manin obstruction over function fields. See for instance [20, Th. B.].

The ideas underlying the proof of Theorem 1.3 are the following. First we notice that we may choose a (necessarily discrete) valuation on K_0 such that all the p^l -th roots of a given infinitely p -divisible point lie in a corresponding maximal unramified extension of the completion of K_0 along the valuation (this is Proposition 2.2). Secondly, we notice that an infinitely p -divisible point on B can be recovered from the images of all its p^l -th roots in the reduction of B modulo the ideal of the valuation (this is Lemma 2.4). This correspondence is Galois equivariant and this implies that an infinitely p -divisible point must have an infinite Galois orbit, if it is not a torsion point. This is a contradiction, because we are dealing with algebraic points (see the end of the proof of Proposition 2.3).

Here is the train of thought underlying the proof of Theorem 1.4. To obtain a contradiction, we suppose that we are given a sequence of non-zero points $P_i \in B(K_0^s)$ such that $p \cdot P_i = P_{i-1}$. We then consider the successive quotients of B by the groups generated by the Galois orbits of the P_i . This gives a sequence of abelian varieties, which is shown to have decreasing modular height (see (2)). Using a fundamental result of Zarhin (see Theorem 3.1), we deduce that infinitely many of the quotients are isomorphic and thus an abelian variety isogenous to B is endowed with a non-trivial endomorphism. This contradicts one of the assumptions of Theorem 1.4.

In the following two sections, we prove the results described above in the same order. The two sections are technically and terminologically independent of each other and can be read in any order.

Acknowledgments. We are very grateful to R. Pink for his detailed comments on an earlier version of this text. Without his input, this text would certainly be a lot less clear. We also thank A. Pillay and F. Benoist for explaining their ideas and conjectures to us and for interesting discussions. Finally, we want to thank J.-F. Voloch for an interesting exchange on and around the problems addressed in this article. Thanks also to B. Poonen for his remarks.

2 Proof of Theorem 1.3

Before describing the proof, we would like to point out the following special case (which is not needed in the proof) :

Lemma 2.1. *If the p -rank of B is 0 then we have $B(K_0^s)^\# \subseteq \text{Tor}(B(K_0^s))$.*

Proof (of the lemma) This is a consequence of the Lang-Néron theorem. The details are left to the reader. \square

Theorem 1.3 will be shown to be a simple consequence of the following two propositions.

To formulate them, let S be the spectrum of a complete noetherian local ring of characteristic p . Let K be the fraction field of S . Let \mathcal{A} be an abelian scheme over S .

Write S^{sh} for the spectrum of the strict henselisation of the ring underlying S and L for the fraction field of S^{sh} . See for instance [16, I, 4, p. 38] for the definition of the strict henselization. We choose once and for all a K -embedding $L \hookrightarrow K^s$.

Proposition 2.2. *Suppose that S is a scheme of characteristic p . Suppose that the p -rank of the fibres of \mathcal{A} is constant. Then we have*

$$(p \cdot \mathcal{A}(K^s)) \cap \mathcal{A}(L) = p \cdot \mathcal{A}(L).$$

The proof of Proposition 2.2 will be given further below.

Proposition 2.3. *Suppose that the residue field of the closed point of S is finite. Then we have $A(L)^\# = \text{Tor}(A(L)^\#)$.*

The proof of Proposition 2.3 will be given further below.

Proof of Theorem 1.3 (assuming Propositions 2.2 & 2.3). Let $x \in B(K_0^s)^\#$. We shall show that x is a torsion point.

We may assume without restriction of generality that $x \in B(K_0)$ (otherwise, we replace K_0 by $K_0(\kappa(x))$). After again possibly replacing K_0 by a finite extension, we may suppose that $B[p](K_0) = B[p](\bar{K}_0)$. Let U be a smooth algebraic curve over \mathbb{F} . We may suppose without restriction of generality that there is an abelian scheme \mathcal{B} over U extending B and that the p -rank of the fibres of \mathcal{B} is constant. Let now u be any closed point of U . Let S be the completion of the local ring of U at u and let $\mathcal{A} := \mathcal{B}_S$. Let K be the fraction field of S and L be the fraction field of the strict henselisation S^{sh} of S , as before.

Now let $x_1, x_2, \dots \in B(K_0^s)$ be elements such that $p \cdot x_l = x_{l-1}$ for all $l \geq 2$ and such that $p \cdot x_1 = x$. Such a sequence exists by assumption. Applying Proposition 2.2 to x and all the x_l successively, we conclude that $x \in \mathcal{A}(L)^\#$. We conclude the proof of Theorem 1.3 by appealing to Proposition 2.3. \square

Proof of Proposition 2.2. Let $N := \ker [p]$ be the kernel of the multiplication by p morphism $[p] : \mathcal{A} \rightarrow \mathcal{A}$. This is a (non-reduced) finite group-scheme over S . There is an exact sequence of finite, flat, S -group schemes

$$0 \rightarrow N_0 \rightarrow N \rightarrow N_{\text{et}} \rightarrow 0 \quad (\%)$$

where N_0 is connected and N_{et} is étale. See [24, (1.4)] for this. Since the unit section of N is open and closed (by the assumption on the p -rank), we see that N_0 is an infinitesimal group-scheme.

Thus the decomposition (%) of N leads to S -isogenies of abelian schemes

$$\mathcal{A} \xrightarrow{\phi} \mathcal{A}_1 \xrightarrow{\mu} \mathcal{A}$$

such that μ is étale, ϕ is purely inseparable and $\mu \circ \phi = [p]$.

Let now $x \in (p \cdot A(K^s)) \cap A(L)$. Let $T \hookrightarrow \mathcal{A}_{S^{\text{sh}}}$ be the closed subscheme arising from the section of $\mathcal{A}_{S^{\text{sh}}}$ over S^{sh} associated to x . We have S^{sh} -morphisms $[p]^*T \rightarrow \mu_{S^{\text{sh}}}^*T \rightarrow T$, where the morphism $[p]^*T \rightarrow \mu_{S^{\text{sh}}}^*T$ is purely inseparable over L and the morphism $\mu_{S^{\text{sh}}}^*T \rightarrow T$ is étale. Furthermore, by construction, there is an

element $y \in \mathcal{A}(K^s)$ such that $y \in ([p]^*T)(K^s)$ and such that $p \cdot y = x$. Let V be the connected component of μ^*T containing the image of $\phi_{S^{\text{sh}}}(y)$. We obtain a sequence of S^{sh} -morphisms

$$\text{Spec } K^s \rightarrow ([p]^*T)_V \rightarrow V \rightarrow T \rightarrow A$$

whose composition is x .

Since S^{sh} is strictly henselian, the morphism $V \rightarrow T$ must be an isomorphism. Let now $v \in V$ be the image of the composition $\text{Spec } K^s \rightarrow ([p]^*T)_V \rightarrow V$. Let $w \in ([p]^*T)_V$ be the image of y . Since the morphism $([p]^*T)_V \rightarrow V$ is purely inseparable, the extension of residue fields $\kappa(w)|\kappa(v)$ is purely inseparable. Hence it must be trivial, since the field extension $\kappa(w)|\kappa(T) = L = \kappa(v)$ is separable, because it is a subextension of the separable algebraic extension $K^s|\kappa(T)$. Hence the extension $\kappa(w)|\kappa(T)$ is trivial.

This shows that we actually have $x \in p \cdot A(L)$, which completes the proof of the proposition. \square

Proof of Proposition 2.3. Let us temporarily use the following notation. Let $R := \Gamma(S, \mathcal{O}_S)$ be the local ring underlying S . Let $\mathfrak{m} \subseteq R$ be the maximal ideal of R and let $k := R/\mathfrak{m}$. We write correspondingly $R^{\text{sh}} := \Gamma(S^{\text{sh}}, \mathcal{O}_{S^{\text{sh}}})$ and $\mathfrak{m}^{\text{sh}} \subseteq R^{\text{sh}}$ for the maximal ideal of R^{sh} . We fix an identification of \bar{k} with $R^{\text{sh}}/\mathfrak{m}^{\text{sh}}$. Let also $\widehat{S^{\text{sh}}}$ be the spectrum of the completion $\widehat{R^{\text{sh}}}$ of R^{sh} along \mathfrak{m}^{sh} and let \widehat{L} be the fraction field of $\widehat{R^{\text{sh}}}$.

Let A_0 be the special fibre of \mathcal{A} , ie the fibre of \mathcal{A} over the unique closed point of S .

Let $I_p(A_0(\bar{k})) := \varprojlim_{l \geq 0} A_0(\bar{k})$, where the transition morphisms are all given by multiplication by p (beware : $I_p(A_0(\bar{k}))$ is not the classical Tate module). So an element of $\varprojlim_{l \geq 0} A_0(\bar{k})$ is given by an inverse system

$$\cdots \rightarrow t_l \rightarrow t_{l-1} \rightarrow \cdots \rightarrow t_0$$

where $t_l \in A_0(\bar{k})$ and $t_{l-1} = p \cdot t_l$ for all $l \geq 0$. Now notice that for any $l \geq 1$, the kernel of the reduction map $\mathcal{A}(R^{\text{sh}}/\mathfrak{m}^{\text{sh},l}) \rightarrow A_0(\bar{k})$ is a commutative group, which is killed by multiplication by p^{l-1} . This follows from the fact that the kernel of the reduction map $\mathcal{A}(\widehat{S^{\text{sh}}}) \rightarrow A_0(\bar{k})$ can be identified with the points of a commutative formal group. See [6, Th. 8.5.9 (a), p. 213].

For a fixed $r \geq 1$, let $\lambda_r : I_p(A_0(\bar{k})) \rightarrow \mathcal{A}(R^{\text{sh}}/\mathfrak{m}^{\text{sh},r})$ be the map sending the inverse system

$$\cdots \rightarrow t_r \rightarrow t_{r-1} \rightarrow \cdots \rightarrow t_0$$

to the element $p^{r-1} \cdot \tilde{t}_{r-1}$, where \tilde{t}_{r-1} is any lifting of t_{r-1} to $\mathcal{A}(R^{\text{sh}}/\mathfrak{m}^{\text{sh},r})$. Notice that the composition of λ_r with the reduction map $\mathcal{A}(R^{\text{sh}}/\mathfrak{m}^{\text{sh},r}) \rightarrow \mathcal{A}(R^{\text{sh}}/\mathfrak{m}^{\text{sh},r-1})$ is the map λ_{r-1} . Hence we obtain a homomorphism

$$\lambda : I_p(A_0(\bar{k})) \rightarrow \varprojlim_r \mathcal{A}(R^{\text{sh}}/\mathfrak{m}^{\text{sh},r}) \simeq \mathcal{A}(\widehat{S^{\text{sh}}}) \simeq \mathcal{A}(\widehat{L}).$$

Remark. A variant of the map λ appears in the theory of the Serre-Tate lifting of ordinary abelian varieties over finite fields (see [12]).

Lemma 2.4. *We have $\mathcal{A}(\widehat{L})^\# = \text{Im}(\lambda)$.*

Lemma 2.4 is a variant of [22, Lemme 3.2.1]. The proofs of both Lemmata are actually identical.

Proof (of the lemma). Let $x_0, x_1, \dots \in A(\widehat{L})$ be such that $p \cdot x_i = x_{i-1}$ for all $i \geq 1$. Let $\rho : A(\widehat{L}) \rightarrow A_0(\bar{k})$ be the reduction map. Consider the inverse system $q \in I_p(A_0(\bar{k}))$ given by

$$\cdots \rightarrow \rho(x_i) \rightarrow \rho(x_{i-1}) \rightarrow \cdots \rightarrow \rho(x_0).$$

We claim that $\lambda(q) = x_0$. For this, it is sufficient to prove that

$$\lambda_r(q) = x_0 \pmod{\mathfrak{m}^{\text{sh},r}}$$

for all $r \geq 1$. We compute

$$\lambda_r(q) = p^{r-1} \cdot (x_{r-1} \pmod{\mathfrak{m}^{\text{sh},r}}) = (p^{r-1} \cdot x_{r-1}) \pmod{\mathfrak{m}^{\text{sh},r}} = x_0 \pmod{\mathfrak{m}^{\text{sh},r}}.$$

This shows that $\cap_{l \geq 0} p^l A(\widehat{L}) \subseteq \text{Im}(\lambda)$.

For the opposite inclusion (which is actually not needed in the text), let

$$\cdots \rightarrow t_1 \rightarrow t_0$$

be an element of $I_p(A_0(\bar{k}))$. This element is the image under multiplication by p of the element

$$\cdots \rightarrow t_2 \rightarrow t_1$$

(in other words $I_p(A_0(\bar{k}))^\# = I_p(A_0(\bar{k}))$). Hence any image by λ of an element of $I_p(A_0(\bar{k}))$ is infinitely p -divisible. \square

Remark. Until now in the proof, we only used the fact that k is a field of characteristic p (not that it is a finite field). In particular, the last lemma is true in that generality.

Now notice that the map λ is compatible with the natural action of $\text{Gal}(\bar{k}|k)$ on $I_p(A_0(\bar{k}))$ and $\mathcal{A}(\widehat{L})$. Let $\sigma \in \text{Gal}(\bar{k}|k) \simeq \widehat{\mathbb{Z}}$ be the Frobenius automorphism. Let $x \in \mathcal{A}(L)^\# \subseteq \mathcal{A}(\widehat{L})^\# = \text{Im}(\lambda)$. Since the residue field of x lies in an algebraic extension of K , there exists $r \geq 1$ such that $\sigma^r(x) = x$. On the other hand, by the Weil conjectures, there is a polynomial Q with integer coefficients and no roots of unity among its roots, such that $Q(\sigma^r) = 0$ on $I_p(A_0(\bar{k}))$. Hence $Q(\sigma^r(x)) = Q(1) \cdot x = 0$. Since $Q(1) \neq 0$, this implies that x is a torsion point. This proves Proposition 2.3. \square

Remark. Let L_1 be the algebraic closure of L in \widehat{L} (notice that if S is a discrete valuation ring then $L_1 = L$; see for instance [13, Ex. 8.3.34, p. 360]). The above proof of Proposition 2.3 actually shows that

$$\mathcal{A}(\widehat{L})^\# \cap \mathcal{A}(L_1) = \text{Tor}(\mathcal{A}(L_1)),$$

which is stronger.

3 Proof of Theorem 1.4

This section can be read independently of the previous one.

To prove Theorem 1.4, we shall crucially need the following theorem. Let U be a smooth proper curve over \mathbb{F} , whose function field is K_0 . A semiabelian scheme \mathcal{A}/U is a commutative group scheme over U , whose geometric fibres are semiabelian varieties. We shall say that a semiabelian scheme over U is generically abelian if its geometric generic fibre is an abelian variety. If \mathcal{A}/U is a semiabelian scheme with zero-section ϵ , we shall write

$$\omega_{\mathcal{A}/U}^0 := \epsilon^*(\det(\Omega_{\mathcal{A}/U}))$$

for the determinant of the sheaf of differentials of \mathcal{A} over U , restricted to the zero-section.

Theorem 3.1 (Zarhin). *Let $\beta, g, n \geq 0$. Suppose that $(n, p) = 1$ and that $n \geq 3$. Up to isomorphism, there is only a finite number of generically abelian semia-
belian schemes \mathcal{A}/U of relative dimension g over U , such that*

- $\deg_U(\omega_{\mathcal{A}/U}^0) = \beta$;
- *there exists an isomorphism of group schemes $\mathcal{A}_{K_0}[n] \simeq (\mathbb{Z}/n\mathbb{Z})_{K_0}^{2g}$.*

The Theorem 3.1 is well-known to specialists but I could find no formal proof of it in the literature.

Proof (of Theorem 3.1). We shall first prove the following statement. Let $\beta \in \mathbb{N}$.

Up to isomorphism, there is only a finite number of generically abelian semia-
belian schemes \mathcal{A}/U of relative dimension g over U , such that (*)

- there exists a principal polarisation on \mathcal{A}_{K_0} ;
- there exists a symplectic isomorphism $(\mathbb{Z}/n\mathbb{Z})_{K_0}^{2g} \simeq \mathcal{A}_{K_0}[n]$;
- $\deg_U(\omega_{\mathcal{A}/U}^0) = \beta$.

To prove this, we shall use the following deep results of D. Mumford, A. Grothendieck, L. Moret-Bailly, C.-L. Chai and G. Faltings.

Let $\mathbf{A}_{g,n}$ be the functor from the category of locally noetherian \mathbb{F}_p -schemes to the category of sets, such that

$$\begin{aligned} \mathbf{A}_{g,n}(S) = \{ & \text{isomorphism classes of the following objects :} \\ & \text{principally polarized abelian schemes over } S \text{ endowed} \\ & \text{with a symplectic isomorphism } (\mathbb{Z}/n\mathbb{Z})_S^{2g} \simeq \mathcal{A}[n] \} \end{aligned}$$

Then D. Mumford proves (see [18]) that the functor $\mathbf{A}_{g,n}$ is representable by a scheme, which is separated and of finite type over \mathbb{F}_p . We shall also denote this scheme by $\mathbf{A}_{g,n}$.

Furthermore, in [5, V, 2., Th. 2.5], C. Chai and G. Faltings prove that there exists

- a scheme $\mathbf{A}_{g,n}^*$, which is proper over \mathbb{F}_p ;

- an open immersion $\mathbf{A}_{g,n} \hookrightarrow \mathbf{A}_{g,n}^*$;
- a semiabelian scheme \mathcal{B} over $\mathbf{A}_{g,n}^*$, such that $\mathcal{B}_{\mathbf{A}_{g,n}}$ is isomorphic to the abelian scheme underlying the universal object over $\mathbf{A}_{g,n}$.

Also they show that $\omega_{\mathcal{B}/\mathbf{A}_{g,n}^*}^0$ is an ample line bundle.

Now write $Z := U \times_{\mathbb{F}_p} \mathbf{A}_{g,n}^*$. Recall that the Hilbert scheme $\mathrm{Hilb}(Z/\mathbb{F}_p)$ is a scheme over \mathbb{F}_p , which is locally of finite type and such that

$$\mathrm{Hilb}(Z/\mathbb{F}_p)(T) = \{\text{closed subschemes of } Z_T, \text{ which are proper and flat over } T\}$$

for any locally noetherian scheme T over \mathbb{F}_p (see [7]).

Furthermore, fix $\Phi \in \mathbb{Q}[\lambda]$, a polynomial with rational coefficients and L/Z an ample line bundle. The \mathbb{F}_p -scheme $\mathrm{Hilb}_\Phi(Z/\mathbb{F}_p)$ is then characterized by the property that

$$\begin{aligned} \mathrm{Hilb}_\Phi(Z/\mathbb{F}_p)(T) := \\ \{\text{closed subschemes } W \text{ of } Z_T, \text{ which are proper and flat over } T \\ \text{and such that } \chi(W_t, L_{W_t}^{\otimes \lambda}) = \Phi(\lambda) \text{ for all } \lambda \in \mathbb{N} \text{ and all } t \in T\} \end{aligned}$$

Here W_t is the fibre at $t \in T$ of the morphism $W \rightarrow T$ and L_{W_t} is the pull-back of L to W_t by the natural morphism $W_t \rightarrow Z$. The symbol $\chi(\cdot)$ refers to the Euler characteristic; by definition

$$\chi(W_t, L_{W_t}^{\otimes \lambda}) = \sum_{r \geq 0} (-1)^r \dim_{\kappa(t)} H^r(W_t, L_{W_t}^{\otimes \lambda}).$$

It is shown in [7], that $\mathrm{Hilb}_\Phi(Z/\mathbb{F}_p)$ is projective over \mathbb{F}_p (as a consequence of the projectivity of Z). Notice that by construction, we have a decomposition

$$\mathrm{Hilb}(Z/\mathbb{F}_p) = \coprod_{\Phi \in \mathbb{Q}[\lambda]} \mathrm{Hilb}_\Phi(Z/\mathbb{F}_p).$$

Finally, it is shown in [6, part II, 5.23] that the functor $\mathrm{Mor}_{\mathbb{F}_p}(U, \mathbf{A}_{g,n}^*)$ from locally noetherian \mathbb{F}_p -schemes T to the category of sets, such that

$$\mathrm{Mor}_{\mathbb{F}_p}(U, \mathbf{A}_{g,n}^*)(T) = \{T\text{-morphisms from } U_T \text{ to } \mathbf{A}_{g,n,T}^*\}$$

is representable by an open subscheme of $\text{Hilb}(Z/\mathbb{F}_p)$. The open immersion $\text{Mor}_{\mathbb{F}_p}(U, \mathbf{A}_{g,n}^*) \hookrightarrow \text{Hilb}(U \times_{\mathbb{F}_p} \mathbf{A}_{g,n}^*/\mathbb{F}_p)$ is described by the natural transformation of functors

$$T\text{-morphism } f \text{ from } U_T \text{ to } \mathbf{A}_{g,n,T}^* \mapsto \text{graph of } f$$

Now fix an ample line bundle M on U . Let L be the line bundle $M \boxtimes \omega_{\mathcal{B}/\mathbf{A}_{g,n}^*}^0$ on $Z = U \times_{\mathbb{F}_p} \mathbf{A}_{g,n}^*$.

Now we are finally ready to tackle our proof of finiteness. Suppose that we are given \mathcal{A}/U as in (*). Restricting to the generic point of U , we get a morphism $\text{Spec } \kappa(U) \rightarrow \mathbf{A}_{g,n}^*$ (whose image is in $\mathbf{A}_{g,n}$) and since $\mathbf{A}_{g,n}^*$ is proper over \mathbb{F}_p and U is a Dedekind scheme, this extends to a morphism $\phi : U \rightarrow \mathbf{A}_{g,n}^*$. We thus get a point $\phi \in \text{Mor}_{\mathbb{F}_p}(U, \mathbf{A}_{g,n}^*)(\mathbb{F}_p)$. Let $\Gamma_\phi \hookrightarrow U \times_{\mathbb{F}_p} \mathbf{A}_{g,n}^*$ be the graph of ϕ . We compute its Hilbert polynomial

$$\begin{aligned} \chi(\Gamma_\phi, L_\Gamma^{\otimes \lambda}) &= \chi(U, (M \otimes \phi^*(\omega_{\mathcal{B}/\mathbf{A}_{g,n}^*}^0))^{\otimes \lambda}) = \deg_U((M \otimes \phi^*(\omega_{\mathcal{B}/\mathbf{A}_{g,n}^*}^0))^{\otimes \lambda}) + 1 - g_U \\ &= \lambda \cdot \deg_U(M \otimes \phi^*(\omega_{\mathcal{B}/\mathbf{A}_{g,n}^*}^0)) + 1 - g_U \\ &= \lambda \cdot \deg_U(M \otimes \omega_{\mathcal{A}/U}) + 1 - g_U \\ &= \lambda \cdot \deg_U(M) + \lambda \cdot \deg_U(\omega_{\mathcal{A}/U}) + 1 - g_U \\ &= \lambda \cdot \deg_U(M) + \lambda \cdot \beta + 1 - g_U := Q(\lambda). \end{aligned} \tag{1}$$

Here g_U is the genus of U . The second equality is justified by the Riemann-Roch theorem on U . From (1) we see that the Hilbert polynomial of $\phi \in \text{Mor}_{\mathbb{F}_p}(U, \mathbf{A}_{g,n}^*)(\mathbb{F}_p)$ only depends on β (once M is given) and thus $\phi \in \text{Hilb}_{Q(\lambda)}(U \times_{\mathbb{F}_p} \mathbf{A}_{g,n}^*/\mathbb{F}_p)(\mathbb{F}_p)$.

Now to prove that there are only a finite number of generically abelian semiabelian scheme \mathcal{A}/U satisfying (*), just notice that the set $\text{Hilb}_{Q(\lambda)}(U \times_{\mathbb{F}_p} \mathbf{A}_{g,n}^*/\mathbb{F}_p)(\mathbb{F}_p)$ is finite, since $\text{Hilb}_{Q(\lambda)}(U \times_{\mathbb{F}_p} \mathbf{A}_{g,n}^*/\mathbb{F}_p)$ is projective and hence of finite type over \mathbb{F}_p .

To conclude, recall the following facts.

First, a generically abelian semiabelian scheme \mathcal{A}/U is completely determined by its generic fibre \mathcal{A}_{K_0} - see [21, IX, cor. 1.4, p. 130]. Secondly, by Zarhin's "trick" (see [17, IX, 1.]), for any abelian variety C/K_0 , the abelian variety $(C \times_{K_0} C^\vee)^4$ can be principally polarized. Thirdly, a given abelian variety C/K_0 has only a finite number of direct factors (see [15, Th. 18.7]). Fourthly, if C/K_0 is an abelian variety, which extends to a semiabelian scheme \mathcal{C} over U , then the dual abelian

variety C^\vee has the same property (by Grothendieck's criterion [9, IX, Prop. 5.13 (c)]) and furthermore $\deg(\omega_{C/U}^0) = \deg(\omega_{C^\vee/U}^0)$ (here we wrote somewhat sloppily C^\vee for the semiabelian extension of C^\vee). See [5, V, 3., Lemma 3.4] for the latter equality.

Putting all these facts together, the theorem readily follows from the just proven fact that there are only a finite number of generically abelian semiabelian scheme \mathcal{A}/U satisfying (*). \square

Proof (of Theorem 1.4). By contradiction. So suppose that there exists a point $P \in B(K_0^s)^\#$ such that $P \neq 0$ and $p^r \cdot P = 0$ for some $r \geq 1$. Then there exists a sequence of points $(P_n \in B(K_0^s))_{n \in \mathbb{N}^*}$ such that $P = P_1$ and $p \cdot P_{i+1} = P_i$ for all $i \in \mathbb{N}^*$. Let G_i be the subgroup of $B(K_0^s)^\#$ generated by the elements $\gamma(P_i)$, where γ runs through $\text{Gal}(K_0^s|K_0)$. By Galois descent, the groups G_i are naturally defined over K_0 and we have natural inclusion morphisms $G_1 \hookrightarrow G_2 \hookrightarrow \dots$ over K_0 . Furthermore, the order d_i of the group G_i is strictly increasing as a function of $i \in \mathbb{N}^*$.

Now we may replace K_0 by a finite extension and U by the corresponding projective curve over k , without restricting generality. Hence we may suppose that B extends to a generically abelian semiabelian scheme \mathcal{B} over U (by Grothendieck's semiabelian reduction theorem [9, IX]). Furthermore, we may suppose that there is an isomorphism $\rho_B : (\mathbb{Z}/l\mathbb{Z})_{K_0}^{2g} \simeq \mathcal{B}_{K_0}[l]$, where $l \geq 3$ is a natural number prime to p . We fix such an l for the rest of the proof.

Now look at the sequence of K_0 -isogenies

$$B \rightarrow B/G_1 \rightarrow B/(G_1 \cdot G_2) \rightarrow \dots B/(G_1 \cdot G_2 \cdot G_3) \rightarrow \dots \quad (2)$$

Let $B_i := B/(G_1 \cdots G_i)$ and $B_0 := B$. Let $\pi_i : \mathcal{B}_i \rightarrow U$ be the connected component of the Néron model of B_i over U . Since all the B_i are isogenous, the criterion [9, IX, Prop. 5.13 (c)] shows that all the \mathcal{B}_i are generically abelian semiabelian schemes over U . Furthermore, by the universal property of Néron models, the morphisms in (2) extend to U -morphisms

$$\mathcal{B} \xrightarrow{\phi_0} \mathcal{B}_1 \xrightarrow{\phi_1} \mathcal{B}_2 \rightarrow \dots \quad (3)$$

Finally, since the isogenies ϕ_i are of degree prime to l , we obtain isomorphisms $\phi_i \circ \phi_{i-1} \circ \dots \circ \phi_0 \circ \rho_B : (\mathbb{Z}/l\mathbb{Z})_{K_0}^{2g} \simeq \mathcal{B}_{i,K_0}[l]$ for all i .

By construction, the morphisms ϕ_i are generically étale. Looking at differentials, we see that there is an exact sequence of coherent sheaves

$$\phi_i^* \Omega_{\mathcal{B}_{i+1}/U} \rightarrow \Omega_{\mathcal{B}_i/U} \rightarrow \Omega_{\phi_i} \rightarrow 0$$

since $\Omega_{\mathcal{B}_{i+1}/U}$ is locally free and since the morphism of sheaves $\phi_i^* \Omega_{\mathcal{B}_{i+1}/U} \rightarrow \Omega_{\mathcal{B}_i/U}$ is injective at the generic point of \mathcal{B}_i , we see that the morphism $\phi_i^* \Omega_{\mathcal{B}_{i+1}/U} \rightarrow \Omega_{\mathcal{B}_i/U}$ is actually injective so that we have an exact sequence

$$0 \rightarrow \phi_i^* \Omega_{\mathcal{B}_{i+1}/U} \rightarrow \Omega_{\mathcal{B}_i/U} \rightarrow \Omega_{\phi_i} \rightarrow 0. \quad (4)$$

Restricting the sequence (4) to the zero section $\epsilon_i : U \rightarrow \mathcal{B}_i$, we get

$$0 \rightarrow \epsilon_i^*(\Omega_{\mathcal{B}_{i+1}/U}) \rightarrow \epsilon_i^*(\Omega_{\mathcal{B}_i/U}) \rightarrow \epsilon_i^*(\Omega_{\phi_i}) \rightarrow 0. \quad (5)$$

Here the exactness on the left is again justified by the fact that the sequence (5) is generically exact on U and the fact that $\epsilon_{i+1}^*(\Omega_{\mathcal{B}_{i+1}/U})$ is locally free. Applying $\deg_U(\cdot)$ to the objects in the last sequence, we conclude that

$$\deg_U(\omega_{\mathcal{B}_i/U}^0) - \deg_U(\epsilon_i^*(\Omega_{\phi_i})) = \deg_U(\omega_{\mathcal{B}_{i+1}/U}^0)$$

Since $\epsilon_i^*(\Omega_{\phi_i})$ is a torsion sheaf, we have $\deg_U(\epsilon_i^*(\Omega_{\phi_i})) \geq 0$ (see [10, Ex. 6.12, chap. 7, p. 149]).

So we conclude that $\deg_U(\omega_{\mathcal{B}_i/U}^0) \leq \deg_U(\omega_{\mathcal{B}/U}^0)$ for all $i \in \mathbb{N}$. From Theorem 3.1, we conclude that there exists $j, l \in \mathbb{N}$ such that $j < l$ and such that $\mathcal{B}_j \simeq \mathcal{B}_l$. Since there is also an étale K_0 -isogeny $B_j \rightarrow B_l$ of degree divisible by p^{l-j} , this means that there is an étale K_0 -isogeny $B_j \rightarrow B_j$ of degree divisible by p^{l-j} . Since this last isogeny cannot coincide with the isogeny given by a power of p (because the isogeny given by multiplication by p always has an inseparable part), this implies that $\text{End}_{\bar{K}_0}(B_j)_{\mathbb{Q}} \neq \mathbb{Q}$. Thus $\text{End}_{\bar{K}_0}(B)_{\mathbb{Q}} \neq \mathbb{Q}$. This concludes the proof. \square

Remark. The Proof of Theorem 1.4 gives a somewhat stronger result. In fact it shows that the conclusion of Theorem 1.4 holds if the following weaker assumption holds : there is no \bar{K}_0 -isogeny $\phi : B_{\bar{K}_0} \rightarrow B'$, such that

- ϕ is étale;
- $\deg(\phi)$ is a power of p ;
- B' carries an étale and finite \bar{K}_0 -endomorphism, whose degree is a power of p .

References

- [1] Dan Abramovich and José Felipe Voloch, *Toward a proof of the Mordell-Lang conjecture in characteristic p* , Internat. Math. Res. Notices **5** (1992), 103–115.
- [2] Franck Benoist, *A theorem of the Kernel in characteristic p* , Preprint. February 8th 2011.
- [3] Franck Benoist, Elisabeth Bouscaren, and Anand Pillay, *Semiabelian varieties over separably closed fields, maximal divisible subgroups, and exact sequences*, Preprint arXiv:0904.2083v1.
- [4] C. Corpet, *Around the Mordell-Lang and Manin-Mumford conjectures in positive characteristic*, in preparation.
- [5] Gerd Faltings and Ching-Li Chai, *Degeneration of abelian varieties*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 22, Springer-Verlag, Berlin, 1990. With an appendix by David Mumford.
- [6] Barbara Fantechi, Lothar Göttsche, Luc Illusie, Steven L. Kleiman, Nitin Nitsure, and Angelo Vistoli, *Fundamental algebraic geometry*, Mathematical Surveys and Monographs, vol. 123, American Mathematical Society, Providence, RI, 2005. Grothendieck’s FGA explained.
- [7] Alexander Grothendieck, *Techniques de construction et théorèmes d’existence en géométrie algébrique. IV. Les schémas de Hilbert*, Séminaire Bourbaki, Vol. 6, Soc. Math. France, Paris, 1995, pp. Exp. No. 221, 249–276 (French).
- [8] *Schémas en groupes. II: Groupes de type multiplicatif, et structure des schémas en groupes généraux*, Séminaire de Géométrie Algébrique du Bois Marie 1962/64 (SGA 3). Dirigé par M. Demazure et A. Grothendieck. Lecture Notes in Mathematics, Vol. 152, Springer-Verlag, Berlin, 1962/1964.
- [9] *Groupes de monodromie en géométrie algébrique. I*, Lecture Notes in Mathematics, Vol. 288, Springer-Verlag, Berlin, 1972. Séminaire de Géométrie Algébrique du Bois-Marie 1967–1969 (SGA 7 I); Dirigé par A. Grothendieck. Avec la collaboration de M. Raynaud et D. S. Rim.
- [10] Robin Hartshorne, *Algebraic geometry*, Springer-Verlag, New York, 1977. Graduate Texts in Mathematics, No. 52.
- [11] Ehud Hrushovski, *The Mordell-Lang conjecture for function fields*, J. Amer. Math. Soc. **9** (1996), no. 3, 667–690.
- [12] Nicholas Katz, *Serre-Tate local moduli*, Algebraic surfaces (Orsay, 1976), Lecture Notes in Math., vol. 868, Springer, Berlin, 1981, pp. 138–202.
- [13] Qing Liu, *Algebraic geometry and arithmetic curves*, Oxford Graduate Texts in Mathematics, vol. 6, Oxford University Press, Oxford, 2002. Translated from the French by Reinie Ern ; Oxford Science Publications.

- [14] Yuri I. Manin, *Letter to the editors: “Rational points on algebraic curves over function fields”* [*Izv. Akad. Nauk SSSR Ser. Mat.* **27** (1963), 1397–1442; MR0157971 (28 #1199)], *Izv. Akad. Nauk SSSR Ser. Mat.* **53** (1989), no. 2, 447–448 (Russian); English transl., *Math. USSR-Izv.* **34** (1990), no. 2, 465–466. MR998307 (90f:11039)
- [15] James S. Milne, *Abelian varieties*, Arithmetic geometry (Storrs, Conn., 1984), Springer, New York, 1986, pp. 103–150.
- [16] ———, *Étale cohomology*, Princeton Mathematical Series, vol. 33, Princeton University Press, Princeton, N.J., 1980.
- [17] Laurent Moret-Bailly, *Pinceaux de variétés abéliennes*, *Astérisque* **129** (1985), 266.
- [18] David Mumford, John Fogarty, and Frances Kirwan, *Geometric invariant theory*, 3rd ed., *Ergebnisse der Mathematik und ihrer Grenzgebiete (2) [Results in Mathematics and Related Areas (2)]*, vol. 34, Springer-Verlag, Berlin, 1994.
- [19] Richard Pink and Damian Roessler, *On ψ -invariant subvarieties of semiabelian varieties and the Manin-Mumford conjecture*, *J. Algebraic Geom.* **13** (2004), no. 4, 771–798.
- [20] Bjorn Poonen and José Felipe Voloch, *The Brauer-Manin obstruction for subvarieties of abelian varieties over function fields*, *Ann. of Math. (2)* **171** (2010), no. 1, 511–532, DOI 10.4007/annals.2010.171.511.
- [21] Michel Raynaud, *Faisceaux amples sur les schémas en groupes et les espaces homogènes*, *Lecture Notes in Mathematics*, Vol. 119, Springer-Verlag, Berlin, 1970 (French).
- [22] ———, *Sous-variétés d’une variété abélienne et points de torsion*, *Arithmetic and geometry*, Vol. I, *Progr. Math.*, vol. 35, Birkhäuser Boston, Boston, MA, 1983, pp. 327–352.
- [23] Damian Rössler, *On the Manin-Mumford and Mordell-Lang conjectures in positive characteristic*, ArXiv 1104.4311. Submitted.
- [24] John Tate, *p -divisible groups.*, *Proc. Conf. Local Fields* (Driebergen, 1966), Springer, Berlin, 1967, pp. 158–183.
- [25] John Tate and José Felipe Voloch, *Linear forms in p -adic roots of unity*, *Internat. Math. Res. Notices* **12** (1996), 589–601.
- [26] José Felipe Voloch, *Diophantine approximation on abelian varieties in characteristic p* , *Amer. J. Math.* **117** (1995), no. 4, 1089–1095, DOI 10.2307/2374961.